

CrossMark
click for updates

Association of immunostaining data with various clinical data and variables of MEST-Oxford classification in IgA nephropathy; a report of our previous study

Hamid Nasri¹

Baradaran Research Laboratory, Isfahan, Iran

Correspondence to:Prof. Hamid Nasri, Email:
hamidnasri@yahoo.com**Received:** 10 May 2024**Accepted:** 23 Sep. 2024**ePublished:** 7 Sep. 2024**Key point**

Our previous study to analyze the relationship between immunostaining data and morphologic variables of the Oxford classification (MEST) in addition to clinical and demographic data in 114 patients with biopsy-proven IgA nephropathy, showed C3 deposition was significantly correlated with serum creatinine levels, while IgA, IgM, and IgG showed no significant relationship.

Keywords: Immunoglobulin A nephropathy, Immunofluorescence, Immunostaining

Citation: Nasri

H. Association of immunostaining data with various clinical data and variables of MEST-Oxford classification in IgA nephropathy; a report of our previous study. *J Prev Epidemiol.* 2025;10(1):e38253. doi: 10.34172/jpe.2025.38253.

**To Editor,**

The analysis for the Oxford classification of IgA nephropathy did not take into account the pattern of immunostaining (1). In 2013, we conducted a study to analyze the relationship between immunostaining data and morphologic variables of the Oxford classification (MEST) in addition to clinical and demographic data in 114 patients with biopsy-proven IgA nephropathy. In our research, we found that only C3 deposition was significantly correlated with serum creatinine levels, while IgA, IgM, and IgG showed no significant relationship. Additionally, our study revealed that IgA deposition scores were positively associated with endocapillary proliferation and segmental glomerulosclerosis in the Oxford classification. Additionally, we observed a positive correlation between the IgM deposition score and the segmental glomerulosclerosis variable. No significant association was found between the IgG deposition score and the four morphologic variables of the Oxford classification. Significant associations were identified between the C3 deposition score and the segmental glomerulosclerosis and endocapillary proliferation variables as well (2). Further research is recommended to explore the aspect of IgA nephropathy and immunostaining data, particularly focusing on the correlation between immunostaining findings and the morphologic variables of the Oxford classification (MEST) as well

as with clinical and demographic data of patients with IgA nephropathy. This exploration could provide valuable insights into the predictive value of immunostaining beyond the established morphologic variables, shedding light on the potential impact of immunostaining patterns on disease prognosis and clinical outcomes in IgA nephropathy (1-3).

Conflicts of interest

The author declares that he has no competing interests.

Ethical issues

Ethical issues (including plagiarism, data fabrication, double publication) have been completely observed by the author.

Funding/Support

None.

References

1. Mubarak M. Significance of immunohistochemical findings in Oxford classification of IgA nephropathy: The need for more validation studies. *J Nephropathol.* 2013;2:210-3. doi: 10.12860/JNP.2013.34.
2. Nasri H, Sajjadih S, Mardani S, Momeni A, Merikhi A, Madihi Y, et al. Correlation of immunostaining findings with demographic data and variables of Oxford classification in IgA nephropathy. *J Nephropathol.* 2013;2:190-5. doi: 10.12860/JNP.2013.30.
3. Catanese L, Rupperecht H, Huber TB, Lindenmeyer MT, Hengel FE, Amann K, Wendt R, Siwy J, Mischak H, Beige J. Non-invasive biomarkers for diagnosis, risk prediction, and therapy guidance of glomerular kidney diseases: a comprehensive review. *Intern J Molecular Sci.* 2024;25:3519.